

NOVEMBER 2020



## Canadian Disability Participation Project

The CDPP is an alliance of university, public, private and government sector partners working together to enhance community participation among Canadians with physical disabilities. The research team for this project has expertise in exercise psychology, spinal cord injury, and disability studies.



### Physical Activity Among Individuals with Spinal Cord Injuries Who Walk

The purpose of this document is to summarize published research that examines physical activity among individuals living with a spinal cord injury (SCI) who have some ability to walk, or ambulate. In particular, we reviewed the amount, types, and outcomes of physical activity in this population, along with the factors that help or hinder them from participating in physical activity.

### People Who Walk with SCI are Overlooked

More than half of SCIs are incomplete, meaning that some recovery of physical function is more likely<sup>1</sup>. Some research suggests that up to 75% of individuals with incomplete SCIs will retain some walking capacity<sup>2,3</sup>, and this number is expected to grow over time<sup>4</sup> as acute care and rehabilitation treatments continue to improve. Yet research continues to focus primarily on individuals with SCI who use a wheelchair, which needs to change if we are to better understand the unique participation experiences of individuals who walk with SCI.

### Physical Activity is Important

No matter whether individuals with SCI walk or use a wheelchair (or both), physical activity is important for maintaining good physical and psychosocial health<sup>5,6</sup>. But very few people with SCI are getting enough physical activity to reap the benefits – and research suggests that people with SCI who walk do half as much leisure-time physical activity than people with SCI who use wheelchairs<sup>7</sup>. Lower levels of leisure-time physical activity might be due to barriers such as greater fatigue and pain<sup>8</sup>, and poorer wheelchair skills<sup>9</sup>. Although it is not known what specific factors help people with SCI who walk to become more active, there is evidence that, like the general population, they need the capability, opportunity, and motivation to be physically active<sup>10</sup>.

We don't really know the types of exercise that are most popular or beneficial for people with SCI who walk. However, one study found that walking is the most frequently reported type of physical activity among 33 ambulatory older adults with a SCI<sup>11</sup>. Another study found that aerobic exercise training can improve cardiovascular health and reduce pain, fatigue, and depression among people with SCI who walk<sup>12</sup>.



## Current Study

Up to this point, there hasn't been systematic research undertaken to examine physical activity among this population. As a first step, we conducted a systematic scoping review in 2020 to identify the amount, types and outcomes of physical activity among individuals with SCI who ambulate<sup>13</sup>, along with factors that prevent them or assist them when participating in physical activity. We wanted to better understand the findings on physical activity in this population to determine what our next steps would be. Synthesizing the research is also an essential step in creating an intervention to support physical activity behaviour for people with SCI who can walk. The review identified 17 articles involving a total of 531 people with SCI who walk. Below is a brief summary.

Category	General Finding	Specific Example
<b>Amount</b>	People with SCI who walk report less activity than wheelchair users with SCI.	Only 7-10% of people who walk with SCI participate in organized sport, compared to 25-28% of wheelchair users with SCI <sup>14,15</sup> .
<b>Type</b>	Studies only looked at leisure-time physical activity (and not other types, like physical activity done at work).	Most studies examined aerobic exercise like treadmill walking <sup>16</sup> or recumbent stepping <sup>12</sup> .
<b>Barriers and Facilitators</b>	Studies examined factors relating to individuals' capability, opportunity, and motivation to do physical activity. The most commonly studied factors were physical capability and reflective motivation.	For reflective motivation, there was a significant positive relationship between attitudes and intentions and physical activity <sup>17</sup> . For physical capability, pain was identified as both a barrier and facilitator for physical activity <sup>17,18</sup> .
<b>Outcomes</b>	Thirteen outcomes related to physical health and one outcome related to psychological health were identified.	A 6-week progressive aerobic exercise program done 3x per week resulted in cardiovascular benefits, more steps taken per day, 20-40% reductions in pain and fatigue, and a 60% decrease in depression <sup>12,19</sup> .

## In Summary

More people are able to walk following SCI than ever before, but those who walk participate in half as much physical activity than people with SCI who use wheelchairs. To improve participation among this population, understanding individuals' capability, opportunity, and motivation to do physical activity is important. We conducted a review of the literature to determine the amount, types, and outcomes of physical activity, as well as the factors that help or hinder participation in physical activity. In addition to confirming that people with SCI who can walk participate in low levels of leisure-time physical activity, our review reveals that most research in this area has examined how individuals' motivation (e.g., attitudes and intentions) and physical capability (e.g., pain, fatigue) influence physical activity. We also know that physical activity is good for improving physical health, but little is known about the impact on other participation outcomes. This review helped us to outline next steps for the research process. These next steps include gaining an understanding the quality of physical activity experiences and developing a physical activity intervention.

## References

1. DeVivo M, Chen Y. Trends in new injuries, prevalent cases, and aging with spinal cord injury. *Arch Phys Med Rehabil.* 2011; 92:332–8.
2. Waters RL, Adkins RH, Yakura JS, Sie I. Motor and sensory recovery following incomplete paraplegia. *Arch Phys Med Rehabil.* 1994;75:67–72.
3. Waters RL, Adkins RH, Yakura JS, Sie I. Motor and sensory recovery following incomplete tetraplegia. *Arch Phys Med Rehabil.* 1994;75:306–11.
4. Nijendijk JHB, Post MWM, Van Asbeck FWA. Epidemiology of traumatic spinal cord injuries in the Netherlands in 2010. *Spinal Cord.* 2014;52:258–63.
5. Williams TL, Smith B, Papathomas A. The barriers, benefits and facilitators of leisure time physical activity among people with spinal cord injury: a meta-synthesis of qualitative findings. *Health Psychol Rev.* 2014;8:404–25.
6. Van der Scheer JW, Martin Ginis KA, Ditor DS, Goosey-Tolfrey VL, Hicks AL, West CR, et al. Effects of exercise on fitness and health of adults with spinal cord injury: a systematic review. *Neurology.* 2017;89:736–45.
7. Martin Ginis KA, Latimer AE, Arbour-Nicitopoulos KP, Buchholz AC, Bray SR, Craven BC, et al. Leisure time physical activity in a population-based sample of people with spinal cord injury part I: demographic and injury-related correlates. *Arch Phys Med Rehabil.* 2010;91:722–8.
8. Saunders LL, Krause JS, DiPiro ND, Kraft S, Brotherton S. Ambulation and complications related to assistive devices after spinal cord injury. *J Spinal Cord Med.* 2013;36:652–9.
9. Bateni H, Maki BE. Assistive devices for balance and mobility: benefits, demands, and adverse consequences. *Arch Phys Med Rehabil.* 2005;86:134–45.
10. Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci.* 2011;6:42.
11. Jörgensen S, Martin Ginis KA, Lexell J. Leisure time physical activity among older adults with long-term spinal cord injury. *Spinal Cord.* 2017;55:848–56.
12. DiPiro ND, Embry AE, Fritz SL, Middleton A, Krause JS, Gregory CM. Effects of aerobic exercise training on fitness and walking-related outcomes in ambulatory individuals with chronic incomplete spinal cord injury. *Spinal Cord.* 2016;54: 675–81.
13. Lawrason SVC, Todd KR, Shaw RB, Martin Ginis KA. Physical activity among individuals with spinal cord injury who ambulate: a systematic scoping review. *Spinal Cord.* 2020;58:735–45.
14. Blauwet C, Sudhakar S, Doherty AL, Garshick E, Zafonte R, Morse LR. Participation in organized sports is positively associated with employment in adults with spinal cord injury. *Am J Phys Med Rehabil.* 2013;92:393–401.
15. Garshick E, Mulroy S, Graves DE, Greenwald K, Horton JA, Morse LR. Active lifestyle is associated with reduced dyspnea and greater life satisfaction in spinal cord injury. *Arch Phys Med Rehabil.* 2016;97:1721–7.
16. Wouda MF, Wejden L, Lundgaard E, Strøm V. Energetic and cardiovascular responses to treadmill walking and stationary cycling in subjects with incomplete spinal cord injury. *Spinal Cord.* 2016;54:51–6.
17. Martin Ginis KA, Papathomas A, Perrier M-J, Smith B. Psychosocial factors associated with physical activity in ambulatory and manual wheelchair users with spinal cord injury: a mixed-methods study. *Disabil Rehabil.* 2017;39:187–92.
18. Robertson T, Bucks RS, Skinner TC, Allison GT, Dunlop SA. Barriers to physical activity in individuals with spinal cord injury: a Western Australian study. *Aust J Rehabil Couns.* 2011;17:74–88.
19. DiPiro MND. Effects of aerobic exercise training on walking and health-related outcomes in individuals with chronic incomplete spinal cord injury [dissertation]. Charleston, SC: Medical University of South Carolina; 2015.

This KT bulletin was prepared by Sarah V. C. Lawrason, PhD Student, UBC.

This work was supported by a Partnership Grant from the Social Sciences and Humanities Research Council of Canada (grant number 895-2013-1021) for the Canadian Disability Participation Project ([www.cdpp.ca](http://www.cdpp.ca)).